

Physicians may offer palliative sedation to unconsciousness to address refractory clinical symptoms, not to respond to existential suffering arising from such issues as death anxiety, isolation, or loss of control. Existential suffering should be addressed through appropriate social, psychological or spiritual support.

*AMA Principles of Medical Ethics: I,VII*

---

### 5.7 Physician-Assisted Suicide

Physician-assisted suicide occurs when a physician facilitates a patient's death by providing the necessary means and/or information to enable the patient to perform the life-ending act (e.g., the physician provides sleeping pills and information about the lethal dose, while aware that the patient may commit suicide).

It is understandable, though tragic, that some patients in extreme duress—such as those suffering from a terminal, painful, debilitating illness—may come to decide that death is preferable to life. However, permitting physicians to engage in assisted suicide would ultimately cause more harm than good.

Physician-assisted suicide is fundamentally incompatible with the physician's role as healer, would be difficult or impossible to control, and would pose serious societal risks.

Instead of engaging in assisted suicide, physicians must aggressively respond to the needs of patients at the end of life. Physicians:

- (a) Should not abandon a patient once it is determined that cure is impossible.
- (b) Must respect patient autonomy.
- (c) Must provide good communication and emotional support.
- (d) Must provide appropriate comfort care and adequate pain control.

*AMA Principles of Medical Ethics: I,IV*

---

### 5.8 Euthanasia

Euthanasia is the administration of a lethal agent by another person to a patient for the purpose of relieving the patient's intolerable and incurable suffering.

It is understandable, though tragic, that some patients in extreme duress—such as those suffering from a terminal, painful, debilitating illness—may come to decide that death is preferable to life.

However, permitting physicians to engage in euthanasia would ultimately cause more harm than good.

Euthanasia is fundamentally incompatible with the physician's role as healer, would be difficult or impossible to control, and would pose serious societal risks. Euthanasia could readily be extended to incompetent patients and other vulnerable populations.

The involvement of physicians in euthanasia heightens the significance of its ethical prohibition. The physician who performs euthanasia assumes unique responsibility for the act of ending the patient's life.