

# NEWS RELEASE



**Date:** Sept. 9, 2010

**Contact:** Christine Stone, Oregon Public Health Information Officer; 971-673-1282, desk; 503-602-8027, cell; [christine.l.stone@state.or.us](mailto:christine.l.stone@state.or.us)

## Rising suicide rate in Oregon reaches higher than national average:

***World Suicide Prevention Day is September 10***

Oregon's suicide rate is 35 percent higher than the national average. The rate is 15.2 suicides per 100,000 people compared to the national rate of 11.3 per 100,000.

After decreasing in the 1990s, suicide rates have been increasing significantly since 2000, according to a new report, "Suicides in Oregon: Trends and Risk Factors," from Oregon Public Health. The report also details recommendations to prevent the number of suicides in Oregon.

"Suicide is one of the most persistent yet preventable public health problems. It is the leading cause of death from injuries – more than even from car crashes. Each year 550 people in Oregon die from suicide and 1,800 people are hospitalized for non-fatal attempts," said Lisa Millet, MPH, principal investigator, and manager of the Injury Prevention and Epidemiology Section, Oregon Public Health.

There are likely many reasons for the state's rising suicide rate, according to Millet. The single most identifiable risk factor associated with suicide is depression. Many people can manage their depression; however, stress and crisis can overwhelm their ability to cope successfully.

Stresses such as from job loss, loss of home, loss of family and friends, life transitions and also the stress veterans can experience returning home from deployment – all increase the likelihood of suicide among those who are already at risk.

"Many people often keep their depression a secret for fear of discrimination. Unfortunately, families, communities, businesses, schools and other institutions often discriminate against people with depression or other mental illness. These people will continue to die needlessly unless they have support and effective community-based mental health care," said Millet.

The report also included the following findings:

- There was a marked increase in suicides among middle-aged women. The number of women between 45 and 64 years of age who died from suicide rose 55 percent between 2000 and 2006 — from 8.2 per 100,000 to 12.8 per 100,000 respectively.