



ETHICS

Physician-Assisted Suicide

Code of Medical Ethics Opinion 5.7

Physician-assisted suicide occurs when a physician facilitates a patient's death by providing the necessary means and/or information to enable the patient to perform the life-ending act (e.g., the physician provides sleeping pills and information about the lethal dose, while aware that the patient may commit suicide).

It is understandable, though tragic, that some patients in extreme duress—such as those suffering from a terminal, painful, debilitating illness—may come to decide that death is preferable to life. However, permitting physicians to engage in assisted suicide would ultimately cause more harm than good.

Physician-assisted suicide is fundamentally incompatible with the physician's role as healer, would be difficult or impossible to control, and would pose serious societal risks.

Instead of engaging in assisted suicide, physicians must aggressively respond to the needs of patients at the end of life. Physicians:

- (a) Should not abandon a patient once it is determined that cure is impossible.
- (b) Must respect patient autonomy.
- (c) Must provide good communication and emotional support.
- (d) Must provide appropriate comfort care and adequate pain control.

AMA Principles of Medical Ethics: I, IV

Read more opinions about this topic

[Code of Medical Ethics: Caring for Patients at the End of Life](#)

Visit the [Ethics main page](#) to access additional Opinions, the Principles of Medical Ethics and more information about the Code of Medical Ethics.